

— ABLYBIO, Help Your Research



# HAGH Rabbit pAb

货号: **AYP21564**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:10 - 1:100
理论分子量	25kDa/28kDa/33kDa
实测分子量	30KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Mitochondrion matrix
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human HAGH (NP_001035517.1).
------	---

## 靶点信息

研究背景	The enzyme encoded by this gene is classified as a thiolesterase and is responsible for the hydrolysis of S-lactoyl-glutathione to reduced glutathione and D-lactate. Three transcript variants encoding different isoforms have been found for this gene.
基因ID	3029
基因名	HAGH
Swiss	Q16775 ( <a href="https://www.uniprot.org/uniprotkb/Q16775/entry">https://www.uniprot.org/uniprotkb/Q16775/entry</a> )
别名	HAGH,GLO2,GLX2,GLXII,HAGH1,HAGH Rabbit pAb,Glyoxalase II

## 产品验证



Western blot analysis of HAGH expressed in HeLa using HAGH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)