

LRRC59 Rabbit pAb

货号: **AYP21565**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:4000
理论分子量	34kDa
实测分子量	35KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Endoplasmic reticulum membrane, Microsome membrane, Nucleus envelope, Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-244 of human LRRC59 (NP_060979.2).
序列	MTKAGSKGGNLRDKLDGNELDLSLSDLNEVPVKELAAALPKATILDLSCKLTTLPSTDFCGLTHLVKLDLSKNKLQQLPADFGRLVNLQHLDLLNNKLVTLPVSAQLKNLKWLDLKDNPVLAQVAGDCLDEKQCKQCANKVLQHMKAQADQERERQRRLEVEREAEEKKREKQRAKEAQERELRKREKAEKERRRKEYDALKAAKREKQKPKKEANQAPKSKSGSRPRKPPPRKHTRS

靶点信息

研究背景	Required for nuclear import of FGF1, but not that of FGF2. Might regulate nuclear import of exogenous FG F1 by facilitating interaction with the nuclear import machinery and by transporting cytosolic FGF1 to, and possibly through, the nuclear pores.
------	---

基因ID	55379
基因名	LRRC59
Swiss	Q96AG4
别名	LRRC59;PRO1855;p34

产品验证



Western blot analysis of LRRC59 expressed in 293T using LRRC59 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn