

TPD52 Rabbit pAb

货号: **AYP21568**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:1000 - 1:2000 IHC: 1:50 - 1:200
理论分子量	16kDa/19kDa/22kDa/24kDa/25kDa/26kDa
实测分子量	23KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	cytoplasm,endoplasmic reticulum,perinuclear region of cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-184 of human TPD52 (NP_005070.1).
序列	MDRGEQGLLRTPVPEEGEDVAATISATETLSEEEQEELRRELAKVEEEIQTL SQVLAKEKHLAEIKRKLGINSLQELKQNI AKGWQDVTATSAYKKTSETLSQAGQKASAAFSSVGSVITKKLEDVKN SPTFKSFEEKVENLKSKVGGTKPAGGDFGEVLNS AANASATTEPLPEKTQESL

靶点信息

研究背景	Enables calcium ion binding activity and protein homodimerization activity. Involved in B cell differentiation. Located in endoplasmic reticulum and perinuclear region of cytoplasm.
基因ID	7163

基因名	TPD52
Swiss	P55327
别名	TPD52;D52;N8L;PC-1;PrLZ;hD52

产品验证



Western blot analysis of TPD52 expressed in HeLa using TPD52 Rabbit pAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn