

GTF3C3 Rabbit pAb

货号: **AYP21624**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	
实测分子量	101kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HeLa,293T,RAW264.7,NIH/3T3,Mouse lung
细胞定位	nuclear membrane,nucleolus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human GTF3C3 (NP_036218.1).
序列	MSGFSPOLIDYLEGKISFEEFERRREERKTREKKSLEKGLSAEENPDDSEVPSSSGINSTKSQDKDVNEGETSDGVRKS VHKVFASMLGENEDDEEEEEEEEEEEEEETPEQPTAGDVFVLEMLNRETMMKEKRPRSKLPRALRGLMGEANIRF

靶点信息

研究背景	The protein encoded by this gene is part of the TFIIC2 complex, which binds to the promoters of small nuclear and cytoplasmic RNA genes in order to recruit RNA polymerase III. The TFIIC2 complex is composed of six subunits. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011]
------	---

基因ID	9330
基因名	GTF3C3
Swiss	Q9Y5Q9
别名	TFIIIC102;TFIIICgamma;TFiiiC2-102;GTF3C3

产品验证



Western blot analysis of GTF3C3 expressed in Mouse lung, HepG2, HeLa using GTF3C3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn