

# TMEM2 Rabbit pAb

货号: **AYP21638**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	200kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung
细胞定位	cytosol,extracellular exosome,nucleolus,plasma membrane
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1150-1250 of human TMEM2 (NP_037522.1).
序列	ERVKIQAATDSKDISNCKMAKAYPQYYRKPSVVKRMPAMLTGLCQCGTRQVVFTSDPHKSYLPVQFQSPDKAETQRGDP SVISVNGTDFTFRSAGVLLLIV

## 靶点信息

研究背景	This gene encodes a type II transmembrane protein that belongs to the interferon-induced transmembrane (IFITM) protein superfamily. The encoded protein functions as a cell surface hyaluronidase that cleaves extracellular high molecular weight hyaluronan into intermediate size fragments before internalization and degradation in the lysosome. It also has an interferon-mediated antiviral function in humans through activation of the JAK/STAT signaling pathway. The activation of this gene by transcription factor SOX4 in breast cancer cells has been shown to mediate the pathological effects of SOX4 on cancer progression. Naturally occurring mutations in this gene are associated with autosomal recessive non-syndromic hearing loss. [provided by RefSeq, Mar 2017]
基因ID	23670
基因名	TMEM2
Swiss	Q9UHN6
别名	TMEM2

## 产品验证



Western blot analysis of TMEM2 expressed in Mouse lung using TMEM2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)