— ABLYBIO, Help Your Research

FADD Rabbit pAb

货号: AYP22353

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	**WB:** 1:500 - 1:1000 **IF/ICC:** 1:50 - 1:200 **IP:** 1:500 - 1:1000
理论分子量	
实测分子量	28KDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse heart
细胞定位	cytoplasm,cytosol,neuron projection,nucleus,plasma membrane,ripoptosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-205 of mouse FADD (NP_034305.1).

靶点信息

研究背景	Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95 or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling.
基因ID	14082
基因名	Fadd
Swiss	Q61160 (https://www.uniprot.org/uniprotkb/Q61160/entry)
别名	Mort1/F,Mort1/FADD,FADD Rabbit pAb,Fadd,FAS-associating death domain-containing protein,Mediator of receptor induced toxicity,Mort1

产品验证



Western blot analysis of FADD expressed in Jurkat,Mouse heart using FADD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)