

DDX55 Rabbit pAb

货号: **AYP22376**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	68kDa
实测分子量	71KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-549,U-87MG,Mouse spleen
细胞定位	cytosol,nucleolus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-360 of human DDX55 (NP_065987.1).
序列	MEHVTEGSWESLPVPLHPQVLGALRELGFYMTVPQSATIPLFMRNKDVAAEAVTGSGKTLAFVIPILEILLRREEKLKKSQV GAIITPTRELAIQIDEVLSHFTKHFPEFSQILWIGGRNPGEDVERFKQGGNIIVATPGRLEDMFRRKAEGLDLASCVRSLDV LVLDEADRLLDMGFEASINTILEFLPKQRRTGLFSATQTQEVENLVRAGLRNPVRVSVKEKGVAASSAQKTPSRLNYYMV CKADEKFNQLVHFLRNHKQEKHLVFFSTCACVEYYGKALEVLVKGVKIMCIHGKMKYKRNKIFMEFRKLQSGILVCTDVMA RGIDIPEVNWVLQYDPPSNASAFVHRCGRTA

靶点信息

研究背景	This gene encodes a member of protein family containing a characteristic Asp-Glu-Ala-Asp (DEAD) motif. These proteins are putative RNA helicases, and may be involved in a range of nuclear processes including translational initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Multiple alternatively spliced transcript variants have been found for this gene. Pseudogenes have been identified on chromosomes 1 and 12.
------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	57696
基因名	DDX55
Swiss	Q8NHQ9
别名	

产品验证



Western blot analysis of DDX55 expressed in Mouse spleen, HeLa, A-549 using DDX55 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn