

LLGL1 Rabbit pAb

货号: **AYP22399**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	112kDa
实测分子量	115KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	OVCAR3,A-431,U-251MG,Mouse testis,Mouse brain,Rat testis
细胞定位	adherens junction,cortical actin cytoskeleton,cytoplasm,cytoskeleton,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human LLGL1 (NP_004131.3).
序列	MMKFRFRRRQGADPQREKLKQELFAFNKTVEHGFPNQPSALAFDPELRIMAIGTRSGAVKIYGAPGVEFTGLHRDAATVTQ MHFLTGGQGRLLSLLDDSSLHLWEIVHHNGCAHLEEALSFQLPSRPGFDGASAPLSLTRVTVLLVAAGDIAALGTEGSSVF FLDVTTLLLEGQTLAPGEVLRVSPDDYR

靶点信息

研究背景	This gene encodes a protein that is similar to a tumor suppressor in Drosophila. The protein is part of a cytoskeletal network and is associated with nonmuscle myosin II heavy chain and a kinase that specifically phosphorylates this protein at serine residues. The gene is located within the Smith-Magenis syndrome region on chromosome 17.
------	---

基因ID	3996
基因名	LLGL1
Swiss	Q15334
别名	DLG4; HUGL; LLGL; Lgl1; Mgl1; HUGL1; HUGL-1

产品验证



Western blot analysis of LLGL1 expressed in Mouse testis,Rat testis,OVCAR3 using LLGL1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn