

# DAB2IP Rabbit pAb

货号: AYP22674

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	131kDa
实测分子量	131kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	axon,cerebellar mossy fiber,climbing fiber,cytoplasm,cytosol,dendrite,parallel fiber,plasma membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 205-320 of human DAB 2IP (NP_619723.1).
序列	RIKARYQTITILPMEMYKEFAEHITNHYLGLCAALEPILSAKTKEEMASALVHILQSTGKVKDFLTDLMMSEVDRCGDNEHLIF RENTLATKAIEEYLKLVGQKYLQDALGEFIKA

## 靶点信息

研究背景	DAB2IP is a Ras (MIM 190020) GTPase-activating protein (GAP) that acts as a tumor suppressor. The DAB 2IP gene is inactivated by methylation in prostate and breast cancers
基因ID	153090

基因名	DAB2IP
Swiss	Q5VWQ8
别名	AIP1; AIP-1; AF9Q34; DIP1/2

## 产品验证



Western blot analysis of DAB2IP expressed in human esophagus cancer, rat heart using DAB2IP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)