

Bcdin3d Rabbit pAb

货号: **AYP22774**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	32kDa
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	cytoplasm,cytosol,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 65-197 of mouse Bcdin3d (NP_083512.2).
------	--

靶点信息

研究背景	O-methyltransferase that specifically monomethylates 5'-monophosphate of cytoplasmic histidyl tRNA (tRNA(His), acting as a capping enzyme by protecting tRNA(His) from cleavage by DICER1. Also able, with less efficiency, to methylate the 5' monophosphate of a subset of pre-miRNAs, acting as a negative regulator of miRNA processing. The 5' monophosphate of pre-miRNAs is recognized by DICER1 and is required for pre-miRNAs processing: methylation at this position reduces the processing of pre-miRNAs by DICER1. Was also reported to mediate dimethylation of pre-miR-145; however dimethylation cannot be reproduced by another group which observes a monomethylation of pre-miR-145.
基因ID	75284

基因名	Bcdin3d
Swiss	Q91YP1
别名	AV138748; 4930556P03Rik

产品验证



Western blot analysis of Bcdin3d expressed in MCF7 using Bcdin3d Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn