

# Pan-Succinyl-K Rabbit pAb

货号: **AYP22880**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	
实测分子量	10-150KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,MCF7,C6,Mouse liver,Mouse heart
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence containing phosphorylated Y.
序列	

## 靶点信息

研究背景	Succinylation of protein is a kind of post-translational modification of protein mainly occurring on lysine residues. Compared with methylation and acetylation, succinylation of lysine has a greater effect on protein structure and function. Sirt5 is an important de-succinylation modifying enzyme involved in the regulation of a variety of biological processes, such as reactive oxygen species defense, fatty acid metabolism and cell apoptosis
基因ID	
基因名	

<b>Swiss</b>	
别名	

## 产品验证



Western blot analysis of Pan-Succinyl-K expressed in Mouse liver,293T,MCF7 using Pan-Succinyl-K Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins : 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)