

— ABLYBIO, Help Your Research



MED26 Rabbit pAb

货号: **AYP22881**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	65kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse brain,Mouse spinal cord
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 500-574 of human MED 26 (NP_004822.2).
------	--

靶点信息

研究背景	The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.
基因ID	9441
基因名	MED26
Swiss	O95402 (https://www.uniprot.org/uniprotkb/O95402/entry)
别名	CRSP7,CRSP70,MED26 Rabbit pAb,MED26,Activator-recruited cofactor 70 kDa component,Cofactor required for Sp1 transcriptional activation subunit 7,Mediator complex subunit 26,Transcriptional coactivator CRSP70,ARC70

产品验证



Western blot analysis of MED26 expressed in Raji, Mouse brain, Mouse spinal cord using MED26 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)