

# RRP8 Rabbit pAb

货号: **AYP22908**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	50kDa
实测分子量	58KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse spleen,Rat kidney,Rat thymus
细胞定位	cytosol,eNoSc complex,nucleolus,nucleoplasm,plasma membrane
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human RRP8 (NP_056139.1).
序列	GPPCSDSEEEVERKKKCHKQALVGSDSAEDKRRKRCQKHAPINSAQHLDNVDQTGPKAWKGSTTNDPPKQSPGSTSP KPPHTLSRKQWRNRQKNKRRCKN

## 靶点信息

研究背景	Essential component of the eNoSC (energy-dependent nucleolar silencing complex, a complex that mediates silencing of rDNA in response to intracellular energy status and acts by recruiting histone-modifying enzymes. The eNoSC complex is able to sense the energy status of cell: upon glucose starvation, elevation of NAD(+)/NADP(+) ratio activates SIRT1, leading to histone H3 deacetylation followed by dimethylation of H3 at 'Lys-9' (H3K9me2 by SUV39H1 and the formation of silent chromatin in the rDNA locus. In the complex, RRP8 binds to H3K9me2 and probably acts as a methyltransferase. Its substrates are however unknown.
------	--

基因ID	23378
基因名	RRP8
Swiss	O43159
别名	NML; KIAA0409

## 产品验证



Western blot analysis of RRP8 expressed in Mouse kidney,Rat kidney,Mouse spleen using RRP8 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)