

TTC7A Rabbit pAb

货号: **AYP22932**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	96kDa
实测分子量	96KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,Rat testis
细胞定位	cytoplasm,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human TTC7A (NP_001275880.1).
序列	MAAKGAHGSYLKVESELERCRAEGHWDRMPQLVRLQTLSPGGGGNRRGSPSAFTFPDTPDDFGKLLLAELLEQCL KENHAKIKDSMPLEKNEPKMSEAKNYLSSILNHGRLSPQYMCEAMLILGKLHYVEGSYRDAISMYARAGIDDMSEKPL YQMRLLSEAFVIKGLSLERLPNSIASRFRLLTEREEVITCFERASWIAQVFLQLEKTTNNSTSRHLKGGCHPLDYELTYFLEAAL QSAYVKNLKKGNIVKGMRELREVLRTVETKATQNFVKVMAAKHLAGVLLHSLSEECY

靶点信息

研究背景	This gene encodes a protein containing tetratricopeptide repeats. Mutations in this gene disrupt intestinal development and can cause early onset inflammatory bowel disease and intestinal atresia. Alternative splicing results in multiple transcript variants.
基因ID	57217

基因名	TTC7A
Swiss	Q9ULT0
别名	TTC7; GIDID; MINAT

产品验证



Western blot analysis of TTC7A expressed in HT-29,Rat testis using TTC7A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn