

CTB4a Rabbit pAb

货号: **AYP22962**

产品信息

| | |
|-------|--|
| 反应 | Oryza sativa |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 120kDa |
| 实测分子量 | 120kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Stems,Spikes |
| 细胞定位 | plasma membrane |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant protein of Oryza sativa CTB4a. |
| 序列 | |

靶点信息

| | |
|------|---|
| 研究背景 | CTB4a (cold tolerance at booting stage), encoding a conserved leucine-rich repeat receptor-like kinase. The different CTB4a alleles confer distinct levels of cold tolerance and selection for variation in the CTB4a promoter region has occurred on the basis of environmental temperature. |
| 基因ID | 4335004 |
| 基因名 | |

| | |
|-------|--------|
| Swiss | Q0JF76 |
| 别名 | CTB4a |

产品验证



Western blot analysis of CTB4a expressed in Stems,Spikes using CTB4a Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn