

# Phospho-AMPKa1-T183/AMPKa2-T172 Rabbit pAb

货号: **AYP23109**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	64kDa/65kDa/62kDa
实测分子量	62KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	apical plasma membrane,cytoplasm,cytosol,nuclear speck,nucleoplasm,nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic phosphorylated peptide around T183 of human AMPKa1/AMPKa2PRKAA1 (NP_006242.5).
序列	LRTSC

## 靶点信息

研究背景	The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.
------	---

基因ID	5562,5563
基因名	PRKAA1,PRKAA2
Swiss	Q13131,P54646
别名	AMPKa1/AMPKa2

## 产品验证



Western blot analysis of Phospho-AMPKa1-T183/AMPKa2-T172 expressed in C6 using Phospho-AMPKa1-T183/AMPKa2-T172 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)