

MYC2 Rabbit pAb

货号: AYP23111

产品信息

反应	Arabidopsis thaliana
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	67kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Seedling,Before inflorescence,Rosette leaf
细胞定位	nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 60-254 of arabidopsis thaliana MYC2 (NP_174541.1).
------	--

靶点信息

研究背景	Encodes a MYC-related transcriptional activator with a typical DNA binding domain of a basic helix-loop-helix leucine zipper motif. Binds to an extended G-Box promoter motif and interacts with Jasmonate ZIM-domain proteins. Its transcription is induced by dehydration stress and ABA treatment. Negative regulator of blue light-mediated photomorphogenic growth and blue and far-red-light-mediated gene expression. Positive regulator of lateral root formation. Regulates diverse JA-dependent functions. Negatively regulates Trp metabolism and biosynthesis of Trp-derived secondary metabolites. Positively regulates flavonoid biosynthesis, resistance to insects, and response to oxidative stress. Regulates other transcription factors, and negatively regulates its own expression.
基因ID	840158

基因名	MYC2
Swiss	Q39204
别名	ATMYC2; F6N18.4; F6N18_4; JAI1; JASMONATE INSENSITIVE 1; JIN1; RD22BP1; ZBF1

产品验证



Western blot analysis of MYC2 expressed in Seedling, Before inflorescence, Rosette leaf using MYC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

实验步骤

访问官网浏览详情: www.ablybio.cn