

IL1RN Rabbit pAb

货号: **AYP23307**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	20kDa
实测分子量	18KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,PC-3
细胞定位	cytoplasm,extracellular region,extracellular space
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-178 of mouse IL1RN (NP_001034790.1).
序列	RPSGKRPCMKQAFRIWDTNQKTFYLRNNQLIAGYLQGPNIKLEEKIDMVPIDLHSVFLGIHGGKLCCLSCAKSGDDIKLQLEE VNITDLSKNKEEDKRFTFIRSEKGPSTSFESAACPGWFLCTTLEADRPVSLTNTPEEPLIVTKFYFQEDQ

靶点信息

研究背景	Predicted to enable cytokine receptor binding activity. Acts upstream of or within insulin secretion and lipid metabolic process. Located in extracellular region and vesicle. Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); eye disease (multiple); kidney failure (multiple); lung disease (multiple); and vascular disease (multiple). Orthologous to human IL1RN (interleukin 1 receptor antagonist).
------	---

基因ID	16181
基因名	
Swiss	P25085
别名	IL-1ra; F630041P17Rik

产品验证



Western blot analysis of IL1RN expressed in A-431,PC-3 using IL1RN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn