

# ARMC10 Rabbit pAb

货号: **AYP23477**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	37KDa
实测分子量	31-37kd
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,MCF7
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 169-308 of human ARM C10 (NP_001154481.1).
序列	TNMTVTNDHQHMLHSYITDLFQVLLTGNGNTKVQVLKLLLNSENAMTEGLLRAQVDSSFLSLYDSHVAKEILLRVLTFLFQNIKNCLKIEGHLAVQPTFTEGSLFFLLHGEECAQKIRALVDHHDAAEVKEKVVTIIPKI

## 靶点信息

研究背景	This gene encodes a protein that contains an armadillo repeat and transmembrane domain. The encoded protein decreases the transcriptional activity of the tumor suppressor protein p53 through direct interaction with the DNA-binding domain of p53, and may play a role in cell growth and survival. Upregulation of this gene may play a role in hepatocellular carcinoma. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 3.
------	--

基因ID	83787
基因名	ARMC10
Swiss	Q8N2F6
别名	SVH; PNAS112; PNAS-112; PSEC0198

## 产品验证



Western blot analysis of ARMC10 expressed in HepG2, MCF7 using ARMC10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)