

HNRNPR Rabbit pAb

货号: **AYP23485**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IHC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IHC: 1:50 - 1:200
理论分子量	59kDa/66kDa/70kDa/71kDa
实测分子量	59kDa/66kDa/70kDa/71kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm,Nucleus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 500 to the C-terminus of human HNRNPR (NP_005817.1).
序列	RGRGGGRGGRGAPPPRGRGAPPPRGRAGYSQRGAPLGPPRGSRGGRGGPAQQQRGRGSRGSRGNRGGNVGGKRRKADGYNQPSKRRQTNNQNWGSQPIAQQPLQQGGDYSGNYGYNNDNQEFYQDTYGGQWK

靶点信息

研究背景	This gene encodes an RNA-binding protein that is a member of the spliceosome C complex, which functions in pre-mRNA processing and transport. The encoded protein also promotes transcription at the c-fos gene. Alternative splicing results in multiple transcript variants. There are pseudogenes for this gene on chromosomes 4, 11, and 10.
------	--

基因ID	10236
基因名	HNRNPR
Swiss	O43390
别名	HNRNPR; HNRPR; hnRNP-R; heterogeneous nuclear ribonucleoprotein R

产品验证



Western blot analysis of HNRNPR expressed in HUVEC using HNRNPR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn