

HMG20B Rabbit pAb

货号: **AYP23489**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	24kDa/32kDa/35kDa
实测分子量	24kDa/32kDa/35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Chromosome, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-317 of human HMG20 B (NP_006330.2).
序列	MSHGPKQPGAAAAPAGGKAPGQHGGFVTVKQEREGEPRAGEKGSHEEPPVKKRGWPKGKKRKKILPNGPKAPVTGY VRFLNERREQIRTRHPDLPFPEITKMLGAEWSKLQPTKQRYLDEAEREKQYMKELRAYQQSEAYKMCTEKIQEKKIKKED SSSGLMNTLLNGHKGGDCDGFSTFDVPIFTEEFLDQNKAREELRRLRKMNVAFEEQNAVLQRHTQSMSSARERLEQEL ALEERTLALQQQLQAVRQALTASFASLPVPGTGETPTLGLDFYMARLHGAIERDPAQHEKLIVRIKEILAQVASEHL

靶点信息

研究背景	Required for correct progression through G2 phase of the cell cycle and entry into mitosis. Required for R COR1/CoREST mediated repression of neuronal specific gene promoters.
基因ID	10362

基因名	HMG20B
Swiss	Q9P0W2
别名	HMG20B; BRAF25; BRAF35; HMGX2; HMGXB2; PP7706; SMARCE1r; SOXL; pp8857; high mobility group 20 B

产品验证



Western blot analysis of HMG20B expressed in human lung cancer using HMG20B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn