

— ABLYBIO, Help Your Research



PSIP1 Rabbit pAb

货号: **AYP23498**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	37kDa/60kDa
实测分子量	37kDa/60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HepG2 cells
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 301-530 of human PSIP 1 (NP_150091.2).
------	--

靶点信息

研究背景	Transcriptional coactivator involved in neuroepithelial stem cell differentiation and neurogenesis. Involved in particular in lens epithelial cell gene regulation and stress responses. May play an important role in lens epithelial to fiber cell terminal differentiation. May play a protective role during stress-induced apoptosis. Isoform 2 is a more general and stronger transcriptional coactivator. Isoform 2 may also act as an adapter to coordinate pre-mRNA splicing. Cellular cofactor for lentiviral integration.
基因ID	11168
基因名	PSIP1
Swiss	O75475 (https://www.uniprot.org/uniprotkb/O75475/entry)
别名	PSIP1,DFS70,LEDGF,PAIP,PSIP2,p52,p75,PC4 and SFRS1-interacting protein,PSIP1 Rabbit pAb,CLL-associated antigen KW-7,Dense fine speckles 70 kDa protein,Lens epithelium-derived growth factor,Transcriptional coactivator p75/p52

产品验证



Western blot analysis of PSIP1 expressed in HepG2 cells using PSIP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)