

NUP210 Rabbit pAb

货号: AYP23525

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	205kDa
实测分子量	250KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	nuclear envelope, nuclear membrane, nuclear pore
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 680-880 of human NUP210 (NP_079199.2).
序列	QNVTAEDTDSIGLALFAPHSSRNYQQHWILVTCQALGEQVIALSVGNKPSLTNPFFAVEPAVVKFVCAPPSRLTLPVYTSPQLDMSCPLLQNKQVVPVSSHNPRLDLAAYDQEGRRFDNFSSLSIQWESTRPVLASIEPELPMQLVVSQDDESGQKKLHGLQAILVHEASGTTAITATATATGYQESHSSARTKQPHDPL

靶点信息

研究背景	The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. The protein encoded by this gene is a membrane-spanning glycoprotein that is a major component of the nuclear pore complex. Multiple pseudogenes related to this gene are located on chromosome 3.
------	---

基因ID	23225
基因名	NUP210
Swiss	Q8TEM1
别名	NUP210; GP210; POM210; nucleoporin 210

产品验证



Western blot analysis of NUP210 expressed in HepG2 using NUP210 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn