

NDUFS5 Rabbit pAb

货号: AYP23572

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	12kDa
实测分子量	12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2 cells, L02 cells, HEK-293 cells, fetal human brain tissue
细胞定位	Mitochondrion,Mitochondrion inner membrane,Mitochondrion intermembrane space,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-106 of human NDUFS 5 (NP_004543.1).
序列	MPFLDIQKRFGNLNIDRWLTIQSGEQPYKMAGRCHAFEKEWIECAHGIGYTRAEKECKIEYDDFVECLLRQKTMRRAGTIRK QRDKLIKEGKYTPPHHIGKGEPRP

靶点信息

研究背景	This gene is a member of the NADH dehydrogenase (ubiquinone) iron-sulfur protein family. The encoded protein is a subunit of the NADH:ubiquinone oxidoreductase (complex I), the first enzyme complex in the electron transport chain located in the inner mitochondrial membrane. Alternative splicing results in multiple transcript variants and pseudogenes have been identified on chromosomes 1, 4 and 17.
------	--

基因ID	4725
基因名	NDUFS5
Swiss	O43920
别名	NDUFS5; CI-15k; CI15K; NADH:ubiquinone oxidoreductase subunit S5

产品验证



Western blot analysis of NDUFS5 expressed in fetal human brain tissue, HepG2 cells, L02 cells using NDUFS5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn