

TDP2 Rabbit pAb

货号: **AYP23620**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:5000
理论分子量	32kDa/40kDa/44kDa
实测分子量	41KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm, Nucleus, PML body, nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human TDP2 (NP_057698.2).
序列	MELGSCLEGGREAAEEEGEPEVKRRLLCVEFASVASCDAAVAQCFLAENDWEMERALNSYFEPPEESALERRPETISEPKTYVDLTNEETTDSTTSKISPEDETQQENGSMFLITWNIDGLDLNNLSERARGVCSYLALYSPDVIFLQEVIPPPSYLKKRSSNYEITGHEEGYFTAIMLKKSRVKLKSQEIIPFP

靶点信息

研究背景	This gene encodes a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor associated factors (TRAFs), and inhibits nuclear factor-kappa-B activation. This protein has sequence and structural similarities with APE1 endonuclease, which is involved in both DNA repair and the activation of transcription factors.
------	---

基因ID	51567
基因名	TDP2
Swiss	O95551
别名	TDP2; AD022; EAP2; EAPII; TTRAP; dj30M3.3; hTDP2; tyrosyl-DNA phosphodiesterase 2

产品验证



Western blot analysis of TDP2 expressed in HeLa using TDP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn