

— ABLYBIO, Help Your Research



CDK2AP1 Rabbit pAb

货号: **AYP23643**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	9kDa/12kDa
实测分子量	9kDa/12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HEK-293 cells, HeLa cells
细胞定位	cytoplasm, cytosol, nucleoplasm, nucleus, perinuclear region of cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-115 of human CDK2A P1 (NP_004633.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
基因ID	8099
基因名	CDK2AP1
Swiss	O14519 (https://www.uniprot.org/uniprotkb/O14519/entry)
别名	CDK2AP1,DOC1,DORC1,ST19,doc-1,p12DOC-1,cyclin-dependent kinase 2-associated protein 1,CDK2AP1 Rabbit pAb,Deleted in oral cancer 1,Putative oral cancer suppressor,CDKAP1

产品验证



Western blot analysis of CDK2AP1 expressed in HEK-293 cells, HeLa cells using CDK2AP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)