

# IGFBP5 Rabbit pAb

货号: **AYP23683**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	30kDa
实测分子量	31KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,U2OS,BT-474
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 21-272 of human IGFBP 5 (NP_000590.1).
序列	LGSFVHCEPCDEKALSMCPPSPLGCELVKEPGCGCCMTCALAEQSCGVYTERCAQGLRCLPRQDEEKPLHALLHGRGVC LNEKSYREQVKIERDSREHEEPTTSEMAEETYSPIFRPKHTRISELKA EAVKKDRRKKLTQSKFVGAENTAHPRIISAPEM RQESEQGPCRRHMEASLQELKASPRMVPRAVYLPNCDRKG FYKRKQCKPSRGRKRGICWCVDKYGMKLPMEYVDGDF QCHTFDSSNVE

## 靶点信息

研究背景	IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.
基因ID	3488

基因名	IGFBP5
Swiss	P24593
别名	IGFBP5; IBP5; insulin-like growth factor-binding protein 5

## 产品验证



Western blot analysis of IGFBP5 expressed in 293T,U2OS,BT-474 using IGFBP5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)