

# TAGLN2 Rabbit pAb

货号: **AYP23740**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	22kDa/24kDa
实测分子量	22kDa/24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH 7.3.
偶联物	Unconjugated
阳性对照	HeLa cells, HEK-293 cells
细胞定位	cytoskeleton, cytosol, extracellular exosome, extracellular region
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-158 of human TAGLN 2 (NP_003555.1).
序列	MANRGPAYGLSREVQQKIEKQYDADLEQILIQWITTQCRKDVGRPQPGRENFQNLKDGTVLCELINALYPEGQAPVKKI QASTMAFKQMEQISQFLQAAERYGINTTDIFQTVDLWEGKNMACVQRTLMLNLGGLAVARDDGLFSGDPNWFPPKSKEN

## 靶点信息

研究背景	The protein encoded by this gene is similar to the protein transgelin, which is one of the earliest markers of differentiated smooth muscle. The specific function of this protein has not yet been determined, although it is thought to be a tumor suppressor. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	8407

基因名	TAGLN2
Swiss	P37802
别名	TAGLN2; HA1756; transgelin-2

## 产品验证



Western blot analysis of TAGLN2 expressed in HeLa cells, HEK-293 cells using TAGLN2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)