

PSRC1 Rabbit pAb

货号: **AYP23749**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	16kDa/33kDa/35kDa/38kDa
实测分子量	16kDa/33kDa/35kDa/38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A431 cells, HEK-293T cells
细胞定位	Cytoplasm,cytoskeleton,spindle,spindle pole
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-215 of human PSRC1 (NP_001005290.1).
序列	MEDLEEDVRFIVDETLDFGGLSPSDSREEEDITVLVTPPEKPLRRGLSHRSDPNAVAPAPQGVRLSLGPLSPEKLEELDEAN RLAAQLEQCALQDRESAGEGLGPRRVKPSPRRETFVLKDSPVRDLLPTVNSLTRSTPSPSSLTPRLRSNDRKGSVRALRAT SGKRPSNMKRESPTCNLFPASKSPASSPLTRSTPPVRGRAGPSGRAAAASEET

靶点信息

研究背景	This gene encodes a proline-rich protein that is a target for regulation by the tumor suppressor protein p53. The encoded protein plays an important role in mitosis by recruiting and regulating microtubule depolymerases that destabilize microtubules. Alternatively spliced transcript variants encoding different isoforms have been described.
基因ID	84722

基因名	PSRC1
Swiss	Q6PGN9
别名	PSRC1; DDA3; FP3214; proline and serine rich coiled-coil 1

产品验证



Western blot analysis of PSRC1 expressed in A431 cells, HEK-293T cells using PSRC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn