

— ABLYBIO, Help Your Research



NR1H2 Rabbit pAb

货号: **AYP23750**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	39kDa/50kDa
实测分子量	63KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 180-260 of human NR1 H2 (NP_009052.4).
------	--

靶点信息

研究背景	The liver X receptors, LXRA (NR1H3; MIM 602423) and LXRβ, form a subfamily of the nuclear receptor superfamily and are key regulators of macrophage function, controlling transcriptional programs involved in lipid homeostasis and inflammation. The inducible LXRA is highly expressed in liver, adrenal gland, intestine, adipose tissue, macrophages, lung, and kidney, whereas LXRβ is ubiquitously expressed. Ligand-activated LXRs form obligate heterodimers with retinoid X receptors (RXRs; see MIM 180245) and regulate expression of target genes containing LXR response elements (summary by Korf et al., 2009 [PubMed 19436111]).[supplied by OMIM, Jan 2010]
基因ID	7376
基因名	NR1H2
Swiss	P55055 (https://www.uniprot.org/uniprotkb/P55055/entry)
别名	NR1H2,LXR-b,LXRβ,NER,NER-I,RIP15,UNR,oxysterols receptor LXR-beta,NR1H2 Rabbit pAb,Liver X receptor beta,Nuclear receptor NER,Nuclear receptor subfamily 1 group H member 2,Ubiquitously-expressed nuclear receptor

产品验证



Western blot analysis of NR1H2 expressed in A-431 using NR1H2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)