

FUNDC1 Rabbit pAb

货号: **AYP23834**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:1000 |
| 理论分子量 | 17kDa |
| 实测分子量 | 17KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | K-562 |
| 细胞定位 | mitochondrial outer membrane |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 50-150 of human FUNDC1 (NP_776155.1). |
| 序列 | VATQIVMGGVTGWCAGFLFQKVGKLAATAVGGGFLLQIASHSGYVQIDWKRVEKDVNKAKRQIKKRANKAAPEINNLIEEATEFIKQNISSGFVGGFL |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes a protein with a FUN14 superfamily domain. The function of the encoded protein is not known. |
| 基因ID | 139341 |
| 基因名 | FUNDC1 |

| | |
|-------|--------|
| Swiss | Q8IVP5 |
| 别名 | FUNDC1 |

产品验证



Western blot analysis of FUNDC1 expressed in K-562 using FUNDC1 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn