

# ECD Rabbit pAb

货号: **AYP24052**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	72kDa/67kDa/76kDa
实测分子量	76KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse brain,Rat brain
细胞定位	cytoplasm,cytosol,nucleoplasm,nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 435-644 of human ECD (NP_009196.1).
序列	ESSEVSKEEKEQNYDLTEVSESMKAFISKVSTHKGAELEPREPSEAPITFDADSFNLYFDKILGPRPNESDSDDLDEDFECL DSDDDLDLFETHEPGEEASLKGTLNLSYMAQMDQELAHTCISKSFTRRNQVEPVSQTTDNNSDEEDSGTGESVMAPV DVDLNLVSNILESYSSQAGLAGPASNLLQSMGVQLPDNTDHRPTSKPTKN

## 靶点信息

研究背景	Enables histone acetyltransferase binding activity. Involved in positive regulation of transcription by RNA polymerase II. Located in cytosol and nucleoplasm.
基因ID	11319

基因名	ECD
Swiss	O95905
别名	GCR2; SGT1; HSGT1

## 产品验证



Western blot analysis of ECD expressed in HeLa, Mouse brain, Rat brain using ECD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)