

— ABLYBIO, Help Your Research



E2A Rabbit pAb

货号: **AYP24255**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	67kDa/68kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Molt-4,Mouse spleen
细胞定位	cytoplasm,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant protein of human E2A.
------	-----------------------------------

靶点信息

研究背景	This gene encodes a member of the E protein (class I) family of helix-loop-helix transcription factors. E proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9.
基因ID	6929
基因名	TCF3
Swiss	P15923 (https://www.uniprot.org/uniprotkb/P15923/entry)
别名	TCF3,AGM8,E2A,E47,ITF1,TCF-3,VDIR,bHLHb21,E2A Rabbit pAb,Class B basic helix-loop-helix protein 21,Immunoglobulin enhancer-binding factor E12/E47,Immunoglobulin transcription factor 1,Kappa-E2-binding factor,Transcription factor 3,Transcription factor ITF-1

产品验证



Western blot analysis of E2A expressed in Molt-4, Mouse spleen using E2A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)