

# CD106 Rabbit pAb

货号: **AYP24796**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	81kDa
实测分子量	85-100KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	RAW264.7(Negative control),Mouse spleen
细胞定位	[Vascular cell adhesion protein 1]: Cell membrane, [Soluble Vascular Cell Adhesion Molecule-1]:
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant protein of mouse CD106.
------	-------------------------------------

## 靶点信息

研究背景	Predicted to enable integrin binding activity and primary amine oxidase activity. Acts upstream of or with in several processes, including cellular response to glucose stimulus; chorio-allantoic fusion; and heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules. Located in apical plasma membrane. Is expressed in several structures, including brain; embryo mesenchyme; extraembryonic component; heart; and hemolymphoid system. Human ortholog(s) of this gene implicated in hypertension. Orthologous to human VCAM1 (vascular cell adhesion molecule 1).
基因ID	22329
基因名	Vcam1

Swiss	P29533
别名	CD106; Vcam-1

## 产品验证



Western blot analysis of CD106 expressed in RAW264.7(Negative control),Mouse spleen using CD106 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)