

CD275 Rabbit pAb

货号: **AYP24820**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	36kDa
实测分子量	65-75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant protein of human CD275.
序列	

靶点信息

研究背景	Predicted to enable identical protein binding activity and signaling receptor binding activity. Involved in positive regulation of interleukin-4 production. Located in external side of plasma membrane. Is integral component of plasma membrane. Is expressed in several structures, including extraembryonic component; gonad; hemolymphoid system gland; liver; and stomach. Orthologous to human ICOSLG (inducible T cell costimulator ligand).
基因ID	50723

基因名	
Swiss	Q9JHJ8
别名	B7h; GI50; GL50; B7-H2; LICOS; B7RP-1; GL50-B; ICOS-L; Icoslg; Ly115l; mKIAA0653

产品验证



Western blot analysis of CD275 expressed in Mouse spleen using CD275 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn