

— ABLYBIO, Help Your Research



Syk Rabbit pAb

货号: **AYP25048**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens IHC: Mus musculus IF: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	69kDa/72kDa
实测分子量	69kDa/72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse spleen,Rat liver
细胞定位	Cell membrane,Cytoplasm,cytosol
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human Syk (NP_001128524.1).
------	---

靶点信息

研究背景	This gene encodes a member of the family of non-receptor type Tyr protein kinases. This protein is widely expressed in hematopoietic cells and is involved in coupling activated immunoreceptors to downstream signaling events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis. It is thought to be a modulator of epithelial cell growth and a potential tumour suppressor in human breast carcinomas. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	6850
基因名	SYK
Swiss	P43405 (https://www.uniprot.org/uniprotkb/P43405/entry)
别名	SYK,p72-Syk,Syk Rabbit pAb,Spleen tyrosine kinase

产品验证



Western blot analysis of Syk expressed in Jurkat, Mouse spleen, Rat liver using Syk Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)