

GSDMD (Full Length+C terminal) Rabbit pAb

货号: **AYP25049**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	53kDa
实测分子量	53KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,THP-1
细胞定位	cytosol,extracellular region,extracellular space,NLRP3 inflammasome complex,nucleoplasm,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 276-484 of human GSDMD (NP_079012.3).
序列	GVPAEAGAFTEDFQGLRAEVETISKELELLDRELCQLLEGLLEGVLRDQLALRALEEALQGGQSLGPVEPLDGPAGAVLECLV LSSGMLVPELAIPVVYLLGALTMLSETQHKLLEAELESQTLLGPLELVGSLLEQSAPWQERSTMSLPPGLLGNWGE GAPAVWLLDECGLELGEDTPHVCWEPQAQGRMCALYASLALLSGLSQEPH

靶点信息

研究背景	Gasdermin D is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. Gasdermin D has been suggested to act as a tumor suppressor. Alternatively spliced transcript variants have been described.
------	--

基因ID	79792
基因名	GSDMD
Swiss	P57764
别名	GSDMD;DF5L;DFNA5L;FKSG10;GSDMDC1

产品验证



Western blot analysis of GSDMD (Full Length+C terminal) expressed in A-431, THP-1 using GSDMD (Full Length+C terminal) Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn