

— ABLYBIO, Help Your Research



Phospho-ULK1-S757 Rabbit pAb

货号: **AYP25065**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus , Homo sapiens , Spodoptera frugiperda IHC: Homo sapiens
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	150kDa
实测分子量	145KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431
细胞定位	Cytoplasm,Preautophagosomal structure,cytosol
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S757 of human Phospho-ULK1-S757 (NP_003556.1).
------	--

靶点信息

研究背景	Serine/threonine-protein kinase involved in autophagy in response to starvation. Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagosomes, the precursors of autophagosomes. Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1 via interaction with RPTOR. Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity. May phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences. Plays a role early in neuronal differentiation and is required for granule cell axon formation. May also phosphorylate SESN2 and SQSTM1 to regulate autophagy. Phosphorylates FLCN, promoting autophagy. Phosphorylates AMBRA1 in response to autophagy induction, releasing AMBRA1 from the cytoskeletal docking site to induce autophagosome nucleation.
基因ID	8408
基因名	ULK1
Swiss	O75385 (https://www.uniprot.org/uniprotkb/O75385/entry)
别名	ULK1,ATG1,ATG1A,UNC51,Unc51.1,hATG1,Phospho-ULK1-S757 Rabbit pAb,Autophagy-related protein 1 homolog,Unc-51-like kinase 1,KIAA0722

产品验证



Western blot analysis of Phospho-ULK1-S757 expressed in A-431 using Phospho-ULK1-S757 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)