

— ABLYBIO, Help Your Research



TRIF / TICAM1 Rabbit pAb

货号: **AYP25083**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IF: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	76kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,U-251MG,22RV1,Mouse spleen,Mouse eye,Rat spleen
细胞定位	Cytoplasmic vesicle,autophagosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-200 of human TRIF / TICAM1 (NP_891549.1).
------	--

靶点信息

研究背景	This gene encodes an adaptor protein containing a Toll/interleukin-1 receptor (TIR) homology domain, which is an intracellular signaling domain that mediates protein-protein interactions between the Toll-like receptors (TLRs) and signal-transduction components. This protein is involved in native immunity against invading pathogens. It specifically interacts with toll-like receptor 3, but not with other TLRs, and this association mediates dsRNA induction of interferon-beta through activation of nuclear factor kappa-B, during an antiviral immune response.
基因ID	148022
基因名	TICAM1
Swiss	Q8IUC6 (https://www.uniprot.org/uniprotkb/Q8IUC6/entry)
别名	IIAE6,MyD88-3,PRVTIRB,TICAM-1,TRIF,TRIF/TICAM1,TICAM1,TRIF / TICAM1 Rabbit pAb,Proline-rich,vinculin and TIR domain-containing protein B,Putative NF-kappa-B-activating protein 502H,Toll-interleukin-1 receptor domain-containing adapter protein inducing interferon beta

产品验证



Western blot analysis of TRIF / TICAM1 expressed in Mouse spleen,Rat spleen,A-549 using TRIF / TICAM1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)