

TRAF3 Rabbit pAb

货号: **AYP25103**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: HeLa
应用	WB IHC IF/ICC
推荐浓度	WB: 1:200 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	55kDa/64kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	22Rv1,SW480,MCF7,U-937,ES-2,Raji,Mouse gastrocnemius muscle,Mouse spleen
细胞定位	Cytoplasm,Endosome,Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human TRAF3 (NP_003291.2).
序列	MESSKKMDSPGALQTNPPLKLHTDRSAGTPVVFVPEQGGYKEKFKVKTVEDKYKCEKCHLVLCSPKQTECGHRFCESMAALLSSSPKCTACQESIVKDKV

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from, members of the TNF receptor (TNFR) superfamily. This protein participates in the signal transduction of CD40, a TNFR family member important for the activation of the immune response. This protein is found to be a critical component of the lymphotoxin-beta receptor (LTbetaR) signaling complex, which induces NF-kappaB activation and cell death initiated by LTbeta ligation. Epstein-Barr virus encoded latent infection membrane protein-1 (LMP1) can interact with this and several other members of the TRAF family, which may be essential for the oncogenic effects of LMP1. Several alternatively spliced transcript variants encoding three distinct isoforms have been reported.
基因ID	7187
基因名	TRAF3
Swiss	Q13114
别名	TRAF3;CAP-1;CAP1;CD40bp;CRAF1;IIAE5;LAP1

产品验证



Western blot analysis of TRAF3 expressed in Mouse gastrocnemius muscle, 22Rv1, SW480 using TRAF3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn