

— ABLYBIO, Help Your Research



Caspase-8 Rabbit pAb

货号: **AYP25128**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Eoma cell , N9 , Mus musculus , Sus scrofa , Ctenopharyngodon idellus , Homo sapiens , Rattus norvegicus , Gallus gallus , Bos taurus , RT-PCR: Ctenopharyngodon idellus IHC: Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	25-32kDa/45- 61kDa
实测分子量	57KDa/18KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C2C12,Jurkat
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 300-400 of human Caspase 8 (NP_203519.1).
------	--

靶点信息

研究背景	This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. This protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined.
基因ID	841
基因名	CASP8
Swiss	Q14790 (https://www.uniprot.org/uniprotkb/Q14790/entry)
别名	ALPS2B,CAP4,Casp-8,FLICE,MACH,MCH5,Caspase 8,CASP8,Caspase-8,Caspase-8 Rabbit pAb,Apoptotic cysteine protease,Apoptotic protease Mch-5,FADD-homologous ICE/ced-3-like protease,FADD-like ICE,ICE-like apoptotic protease 5,MORT1-associated ced-3 homolog

产品验证



Western blot analysis of Caspase-8 expressed in C2C12, Jurkat using Caspase-8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)