

— ABLYBIO, Help Your Research



cIAP2 Rabbit pAb

货号: **AYP25170**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Homo sapiens WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	68kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,RAW264.7
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence within amino acids 75-155 of human cIAP2 (NP_001156.1).
------	--

靶点信息

研究背景	This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified.
基因ID	330
基因名	BIRC3
Swiss	Q13489 (https://www.uniprot.org/uniprotkb/Q13489/entry)
别名	BIRC3,AIP1,API2,CIAP2,HAIP1,HIAP1,MALT2,MIHC,RNF49,c-IAP2,cIAP2 Rabbit pAb,Apoptosis inhibitor 2,Cellular inhibitor of apoptosis 2,IAP homolog C,Inhibitor of apoptosis protein 1,RING finger protein 49,RING-type E3 ubiquitin transferase BIRC3,TNFR2-TRAF-signaling complex protein 1

产品验证



Western blot analysis of cIAP2 expressed in HepG2,RAW264.7 using cIAP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)