

Phospho-BRCA1-S988 Rabbit pAb

货号: AYP25217

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	IF/ICC
推荐浓度	IF/ICC: 1:50 - 1:200
理论分子量	7kDa/78-85kDa/202-210kDa
实测分子量	7kDa/78-85kDa/202-210kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF-7 cells
细胞定位	Chromosome, Cytoplasm, Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息

抗原信息	A synthetic phosphorylated peptide around S988 of human BRCA1 (NP_009225.1).
序列	IKSFV

靶点信息

研究背景	This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast and ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiological function of this gene. Many alternatively spliced transcript variants, some of which are disease-associated mutations, have been described for this gene, but the full-length natures of only some of these variants has been described. A related pseudogene, which is also located on chromosome 17, has been identified.
基因ID	672
基因名	BRCA1
Swiss	P38398
别名	BRCA1;BRCAI;BRCC1;BROVCA1;FANCS;IRIS;PNCA4;PPP1R53;PSCP;RNF53

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn