

HAUS8 Rabbit pAb

货号: **AYP25230**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	38kDa/44kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,NCI-H460,Mouse testis
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center,spindle,spindle pole
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 166-410 of human HAU S8 (NP_219485.1).
序列	NLAEFERRAEKNLLIMCKEKEKLQKKAHELKRRLLLSQRKRELADVLDLDAQIEMLSPFQAVATRFKEQYRTFATALDTRHELP VRSIHLEGGDQQLLDALQHELVTTRQLLGELDVGDSEENVQVLDLLSELKDVTAKKDLELRRSFAQVLELSAEASKEAALA NQEVWEETQGMAPPSRWYFNQDSACRESGGAPKNTPLSEDDNPGASSAPAQATFISPSSEDFSSSSQAEVPPSLRSRSGR DLS

靶点信息

研究背景	HAUS8 is 1 of 8 subunits of the 390-kD human augmin complex, or HAUS complex. The augmin complex was first identified in Drosophila, and its name comes from the Latin verb 'augmentare,' meaning 'to increase.' The augmin complex is a microtubule-binding complex involved in microtubule generation within the mitotic spindle and is vital to mitotic spindle assembly (Goshima et al., 2008 [PubMed 18443220]; Uehara et al., 2009 [PubMed 19369198]).[supplied by OMIM, Jun 2010]
基因ID	93323
基因名	HAUS8
Swiss	Q9BT25
别名	HAUS8;DGT4;HICE1;NY-SAR-48

产品验证



Western blot analysis of HAUS8 expressed in Jurkat, NCI-H460, Mouse testis using HAUS8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn