

# HADHA Rabbit pAb

货号: **AYP25260**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> 293T
应用	<a href="#">WB</a> <a href="#">IF/ICC</a> <a href="#">IP</a>
推荐浓度	<b>WB:</b> 1:2000 - 1:6000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:50 - 1:100
理论分子量	28kDa/83kDa
实测分子量	78KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,K-562
细胞定位	Mitochondrion
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 484-763 of human HADHA (NP_000173.2).
序列	IAAVSKRPEKVI GMHYFSPVDKMQLL EITTEKTSKDTSASAVAVGLKQ GKVIIVVKDGP GFYTTTRCLAPMMSEVIRILQEGVDPKKLDSLTT SFGFPVGAATLVDEVGVDVAKHVAEDLGKVFGERFGG GNPPELLTQMVSKGFLGRKSGKGFYIYQEGVKRKDLNSDMDSILASLKLPPKSEVSSDEDIQFRLVTRFVNEAVMCLQEGILATPAEGDIGAVFGLGFPPCLGGPFRFVDLYGAQKIVDRLKKYEAA YGKQFTPCQLLADHANS PNKKFYQ

## 靶点信息

研究背景	This gene encodes the alpha subunit of the mitochondrial trifunctional protein, which catalyzes the last three steps of mitochondrial beta-oxidation of long chain fatty acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four beta subunits, with the alpha subunit catalyzing the 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratase activities. Mutations in this gene result in trifunctional protein deficiency or LCHAD deficiency. The genes of the alpha and beta subunits of the mitochondrial trifunctional protein are located adjacent to each other in the human genome in a head-to-head orientation.
基因ID	3030
基因名	HADHA
Swiss	P40939
别名	HADHA;ECHA;GBP;HADH;LCEH;LCHAD;MTPA;TP-ALPHA

## 产品验证



Western blot analysis of HADHA expressed in 293T, K-562 using HADHA Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)