

PNKD Rabbit pAb

货号: **AYP25297**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	15kDa/36kDa/40kDa/42kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,22Rv1,THP-1,Mouse brain,Mouse kidney
细胞定位	Cytoplasm,Membrane,Mitochondrion,Nucleus,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human PNKD (NP_056303.3).
序列	MAAVVAATALKGRGARNARVLRGILAGATANKASHNRTRALQSHSSPEGKEEPEPLSPELEYIPRKRGNPKMKAVGLAWY SLYTRTWLGYLFYRQQLRRARNRYPKGHSKTQPRLFNGVKVLPVLSNDNYSYLIIDTQAQLAVAVDPSDPRAVQASIEKEG VTLVAILCTHKHWDHSGGNRDLRRHRDCRVYGPQDGIPYLTHPLCHQDVVSVGRLQIRALATPGHTQGHLVYLLDGEF YKGPSCFLSGDLLFLSGCGRTFEGNAET

靶点信息

研究背景	This gene is thought to play a role in the regulation of myofibrillogenesis. Mutations in this gene have been associated with the movement disorder paroxysmal non-kinesigenic dyskinesia. Alternative splicing results in multiple transcript variants.
------	--

基因ID	25953
基因名	PNKD
Swiss	Q8N490
别名	PNKD;BRP17;DYT8;FKSG19;FPD1;KIPP1184;MR-1;MR1;PDC;PKND1;TAHCCP2

产品验证



Western blot analysis of PNKD expressed in Mouse brain,A-549,22Rv1 using PNKD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn