

— ABLYBIO, Help Your Research



HSD17B3 Rabbit pAb

货号: **AYP25336**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	29kDa/34kDa
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	endoplasmic reticulum
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-310 of human HSD17 B3 (NP_000188.1).
------	--

靶点信息

研究背景	This isoform of 17 beta-hydroxysteroid dehydrogenase is expressed predominantly in the testis and catalyzes the conversion of androstenedione to testosterone. It preferentially uses NADP as cofactor. Deficiency can result in male pseudohermaphroditism with gynecomastia.
基因ID	3293
基因名	HSD17B3
Swiss	P37058 (https://www.uniprot.org/uniprotkb/P37058/entry)
别名	HSD17B3,EDH17B3,SDR12C2,HSD17B3 Rabbit pAb,Estradiol 17-beta-dehydrogenase 2,Short chain dehydrogenase/reductase family 12C member 2,Testicular 17-beta-hydroxysteroid dehydrogenase,Testosterone 17-beta-dehydrogenase 3

产品验证



Western blot analysis of HSD17B3 expressed in 293T using HSD17B3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)