

TUBGCP3 Rabbit pAb

货号: **AYP25349**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	48kDa/93kDa/103kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse spleen,Mouse thymus,Rat testis
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human TUBGCP3 (NP_001273207.1).
序列	MATPDQKSPNVLLQNLCCRILGRSEADVAQQFQYAVRVIGSNFAPTVERDEFVLVAEKIKKELIRQRREADAALFSELHRKLSQGVLNKWSILYLLLSLSEDP RRQPSKVSSYATLFAQALPRDAHSTPYYYARPQTLPLSYQDRSAQSAQSSGSGVSSGISIGLICALSGPAPAPQSLLPGQSNQAPGVGDCLRQQLGSR LAWTLTANQPSSQATTSKGVPSAVSRNMTRSRRREGDTGGTMEITEAAL

靶点信息

研究背景	Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.
基因ID	10426
基因名	TUBGCP3

Swiss	Q96CW5
别名	TUBGCP3;104p;GCP3;Grip104;SPBC98;Spc98;Spc98p

产品验证



Western blot analysis of TUBGCP3 expressed in Mouse brain,Rat testis,Mouse spleen using TUBGCP3 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn