

NIN Rabbit pAb

货号: **AYP25383**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	147kDa/160kDa/236-248kDa
实测分子量	243kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse skin,Rat skin
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 380-665 of human NIN (NP_891991.1).
序列	VREKEKLRSDLDKAEKLSLMASEVDDHHAAIERRNEYNLRLKLDDEYKERIAALKNELRKEREQILQQAGKQRLELEQEIEKA KTEENYIRDRLALSLENENELLENNAEKLAEYENLTNKLQRNLENVLAKEFGDLPSSAEFFLQEERLTQMRNEYERQCR VLQDQVDELQSELEEYRAQGRVLRRLPLKNPSEEEVANSGGIEPEHGLGSEECNPLNMSIEAELVIEQMKEQHHRDICLRL ELEDKVRHYEKQLDETVVSCKAQENMKQRHENETHTL

靶点信息

研究背景	This gene encodes one of the proteins important for centrosomal function. This protein is important for positioning and anchoring the microtubules minus-ends in epithelial cells. Localization of this protein to the centrosome requires three leucine zippers in the central coiled-coil domain. Multiple alternatively spliced transcript variants that encode different isoforms have been reported.
------	---

基因ID	51199
基因名	NIN
Swiss	Q8N4C6
别名	NIN;SCKL7;ninein

产品验证



Western blot analysis of NIN expressed in Mouse skin,Rat skin using NIN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn