

ABCF2 Rabbit pAb

货号: **AYP25385**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	71kDa/72kDa
实测分子量	71KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human ABCF2 (NP_005683.2).
序列	MPSDLAKKKAACKKEAAKARQRPRKGHEENGDVVTEPQVAEKNEANGRETTEVDLLTKELED FEMKKAARAVTGVLAS HPNSTDVHIINLSLTFHGQELLSDTKLELNSGRRYGLIGLNGIGKSMLLSAIGKREVPPIEHIDIYHLTREMPPSDKTPLHCVM EVDTERAMLEKEAERLAHEDAECLELMELYERLEELDADKAEMRASRILHGLGFTPAMQRKKLKDFSGGWRMRVALARAL FIRPFML

靶点信息

研究背景	This gene encodes a member of the ATP-binding cassette (ABC) transporter superfamily. ATP-binding cassette proteins transport various molecules across extra- and intracellular membranes. Alterations in this gene may be involved in cancer progression. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 3 and 7.
------	---

基因ID	10061
基因名	ABCF2
Swiss	Q9UG63
别名	ABCF2;ABC28;EST133090;HUSSY-18;HUSSY18

产品验证



Western blot analysis of ABCF2 expressed in HeLa using ABCF2 Rabbit pAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn